

Surface-active dust in a plasma sheath

A.M. Ignatov,^{1,*} P.P.J.M. Schram,² and S.A. Trigger³

¹*General Physics Institute, Moscow, Russia*

²*Eindhoven University of Technology, Eindhoven, The Netherlands*

³*Humboldt University, Berlin, Germany*

(Dated: February 2, 2008)

Abstract

The inhomogeneity of the plasma pressure near a conducting electrode is a cause for introducing the surface tension associated with the plasma-electrode interface. We evaluate the dependence of the surface tension on the density of the charged dust immersed in the plasma sheath. In a wide range of parameters, the surface tension turns out to be an increasing function of the dust density.

PACS numbers: 52.25.Zb, 52.40.Hf, 52.25.Wz

*Electronic address: aign@fpl.gpi.ru

Throughout the recent decade, an admixture of charged macro-particles, electrons, ions, and neutral atoms, called a dusty plasma, has been the subject to some thousands of studies. Numerous aspects of the physics of dusty plasmas are discussed in the series of reviews starting with [1] and the monograph [2].

Many laboratory experiments deal with relatively small number of dust grains levitating above a horizontal rf-powered or dc-biased electrode. Under these conditions, dust grains are negatively charged and the gravity force is compensated by the strong electric field in a plasma sheath adjacent to an electrode. Dust suspended in a plasma sheath self-organizes itself in various single or multi-layered structures. In particular, recent experiments [3] demonstrated that with growing rf power or gas pressure there evolves a void, that is, a dust-free region, in the center of a single dust layer. Since negatively charged grains levitating at the same height repulse one another via the screened Coulomb potential, there are no evident reasons for the void emergence. The explanation of this phenomenon given in [3] implies existence of an attractive interaction between grains; however, the nature of the interaction remains obscure.

To avoid confusion, it should be stressed that three-dimensional voids, which have been observed earlier [4, 5, 6, 7], evolve in the plasma bulk. The theory of three-dimensional voids [8, 9] demands large number of grains and strong influence of the dust upon the discharge structure. At first sight, both assumptions fail under the conditions of the experiment [3], where some hundreds of grains only have been used.

The main purpose of this Letter is to introduce the concept of plasma surface tension. We demonstrate that even a small number of dust grains can modify the plasma sheath profile in such a way that the sheath may confine the dust in the horizontal direction. Our reasoning is based on the analysis of the plasma pressure in the sheath-presheath area of a gas discharge. We show that the horizontal electric force appearing due to the inhomogeneity of a dust layer is conveniently treated in terms of the plasma surface tension. It should be stressed that explanation of the experiment [3] is beside our purpose: there is evidently a number of relevant factors that are not discussed here.

The schematic of the model implemented here is depicted in Fig. 1. The plasma consisting of electrons and singly charged ions is described with the help the continuity equation

$$\frac{dn_i(z)v_i(z)}{dz} = \nu_{ion}n_e(z), \quad (1)$$




FIG. 1: Schematic of the model

the momentum balance equation

$$\frac{dm_i n_i(z) v_i^2(z)}{dz} + e n_i(z) \frac{d\phi(z)}{dz} = 0, \quad (2)$$

and Poisson's equation

$$\frac{d^2\phi(z)}{dz^2} = 4\pi e(n_e(z) - n_i(z)) - 4\pi\rho_d(z). \quad (3)$$

Here, ν_{ion} is the electron impact ionization frequency and $\rho_d(z)$ is the charge density of aerosol grains. Other notation is standard. Electrons are supposed to be Boltzmannian: $n_e(z) = n_0 \exp(e\phi(z)/T_e)$, where n_0 is a normalizing constant. We also assume that the mean free path of the ion-neutral collisions is large compared to the Debye length, which determines the spatial scale of the sheath.

Since the electron and ion fluxes at the electrode ($z = 0$) are equal, one of the boundary conditions to Eqs. (1-3) is

$$n_i(0)v_i(0) + n_e(0)\sqrt{\frac{T_e}{2\pi m_e}} = 0. \quad (4)$$

Far from the electrode the plasma is quasi-neutral, *i.e.*, $n_e = n_i = n_0$. Generally, the equilibrium plasma density, n_0 , is the eigenvalue of the problem and it is determined by the balance of ionization and losses of charged particles. For this reason, the bulk plasma density may depend on the dust charge located in the sheath because the latter modifies the electric field profile, which, in its turn, influences the net particle flux at the electrode. However, here we assume the overall size of the plasma to be large enough, so that the equilibrium density, n_0 , is conditioned by the recombination in the plasma volume. Therefore, the plasma density

tends to a certain fixed value at $z \rightarrow \infty$. In order to patch the quasi-neutral plasma and the sheath, the equations (1,2) are solved assuming that $n_e(z) = n_i(z)$ and $\rho_d(z) = 0$ (*e.g.*, [10]) that results in

$$n_i(z) = \frac{n_0 v_s^2}{v_s^2 + v_i(z)^2}, \quad (5)$$

where $v_s = \sqrt{T_e/m_i}$ is the ion sound velocity. It is assumed that at the end of the simulation area, $z = z_0$, the ion flow velocity is fixed, *i.e.*, $v_i(z_0) = v_0$. The value of v_0 should be small enough; all solutions exemplified below are obtained for $v_0 = -0.01v_s$. Then, Eqs. (1,5) are used to evaluate boundary values of the density, $n_i(z_0)$, the potential, $\phi(z_0)$, and its derivative, $\phi'(z_0)$; this results in four boundary conditions for Eqs. (1-3). The length of the simulation area, z_0 , is determined with the help of the fifth boundary condition (4).

In the framework of our model, the dust is represented as an infinitesimally thin charged massive layer levitating at $z = z_l$, that is, its charge density is

$$\rho_d(z) = -q\sigma\delta(z - z_l), \quad (6)$$

where $-q$ ($q > 0$) is the charge of a single grain and σ is the surface dust density. The equilibrium levitation height, z_l , depends on the vertical force balance. The largest forces acting upon dust in a plasma sheath are the electric field force and the gravity force. Since the z -component of the electric field, $E(z) = -\phi'(z)$, is discontinuous at $z = z_l$, the vertical force balance is written as

$$\frac{1}{2}q [E(z_l + 0) + E(z_l - 0)] = Mg, \quad (7)$$

where M is the mass of a single grain and g is the gravity acceleration (in Fig. 1, the Earth is at the bottom). On the other hand, the electric field discontinuity is given by

$$E(z_l + 0) - E(z_l - 0) = -4\pi q\sigma \quad (8)$$

and the potential, $\phi(z)$, is a continuous function at $z = z_l$. The relations (7,8) allows us to patch the numeric solutions of Eqs. (1-3) below ($0 < z < z_l$) and above ($z_l < z < z_0$) the layer, and to obtain the equilibrium levitation height, z_l , and the net thickness of the sheath-presheath area, z_0 .

It is well-known that the grain charge, q , in Eqs. (7,8) depends on the plasma parameters, and its variability should be also taken into account [11]. However, here we assume that the grain charge is fixed; the results of preliminary computer runs with the variable grain charge did not alter our main conclusions. We also ignore the ion wind force in Eq. (7) as long as absorption of plasma by dust grains. Simple estimations show that these processes are of minor importance.

Our main objective is to analyze the plasma pressure profile in the sheath-presheath area. Evidently, Eqs. (1-3) ensure the conservation of the net linear momentum of the system. The tensor of the momentum flux, or the pressure tensor is a sum of the ion momentum flux, the electron pressure and Maxwell's stress: $P_{\alpha\beta} = m_i n_i v_{i\alpha} v_{i\beta} + n_e T_e \delta_{\alpha\beta} + \mathbf{E}^2 \delta_{\alpha\beta} / 8\pi - E_\alpha E_\beta / 4\pi$ ($\alpha, \beta = x, y, z$). In the one-dimensional model described above, the pressure tensor is diagonal: $P_{zz} = P_l$ and $P_{xx} = P_{yy} = P_{tr}$. The longitudinal pressure is

$$P_l(z) = m_i n_i(z) v_i(z)^2 + n_e(z) T_e - \frac{1}{8\pi} E(z)^2. \quad (9)$$

Since within our model there are no momentum losses in the plasma, P_l is constant in the absence of dust and it is equal to the plasma electron pressure at $z \rightarrow \infty$, $P_l = n_0 T_e$.

The transverse pressure is written as

$$P_{tr}(z) = n_e(z) T_e + \frac{1}{8\pi} E(z)^2. \quad (10)$$

In contrast with Eq. (9), there is no contribution of the ion momentum flux and the electric field pressure is of the opposite sign.

Although the transverse pressure tends to the electron pressure in the plasma bulk, $P_{tr}|_{z \rightarrow \infty} = n_0 T_e$, it is essentially nonuniform in the sheath-presheath area adjacent to the electrode. This circumstance allows us to define the *surface tension* at the plasma-electrode interface as

$$s(\sigma) = \int_0^\infty dz (n_0 T_e - P_{tr}(z)). \quad (11)$$

The physical meaning of this quantity is fairly evident. Since the bulk plasma pressure is positive, two parts of the discharge repulse one another. As we shall demonstrate later, the transverse pressure (10) is reduced near the wall; therefore, there arises the additional horizontal force attracting two parts of the plasma-wall interface. The surface tension (11)

characterizes the magnitude of this force per unit length. It should be noted that usually surface tension is defined in terms of surface free energy. The latter is obviously meaningless in application to an open, nonequilibrium system like the plasma-wall boundary. However, due to the momentum conservation, the surface tension given by (11) is a well-defined concept. Of particular interest for our purposes is the dependence of the surface tension (11) on the dust density.

The utility of the notion (11) may be illustrated in the following way. Suppose that the layer density is a weakly varying function of the transverse coordinate, x , that is, $\partial\sigma/\partial x \sim \epsilon$, where $\epsilon \ll 1$. Integrating P_{xx} component of the pressure tensor with respect to z at nearby points x and $x + \Delta x$ we see that the horizontal force (*i.e.*, the momentum flux) acting upon a small part of the layer of the width Δx is written in two identical forms:

$$F_x = -q\sigma E_x \Delta x \equiv -\Delta x \frac{\partial}{\partial x} \int dz P_{xx}(x, z). \quad (12)$$

Comparing both parts of the identity and noting that $E_x \sim \partial/\partial x \sim O(\epsilon)$ we conclude that in order to obtain the first-order term of the expansion of the horizontal force in powers of ϵ it is sufficient to evaluate the surface tension (11) ignoring the spatial inhomogeneity of the layer density. Therefore, $f_x = \partial s(\sigma(x))/\partial x$. This simple reasoning may be confirmed by the more tedious asymptotic expansion of the hydrodynamic equations in powers of ϵ .

We can also extract some useful information about the horizontal force even for the sharply bounded layer. Suppose that the dust layer occupies the area $x < 0$, as depicted in Fig. 1. The levitation height, z_l , now depends on x , the electric field and the ion flow near the edge of the layer are essentially two-dimensional. However, the horizontal component of the force is the difference $f_x = s_0 - s_1$, where $s_{0,1}$ are given by the integral (11) evaluated across the sheath in the dust-free area, at $x = x_0$, and in the presence of dust, $x = x_1$ (Fig. 1). Assuming that both points $x_{0,1}$ are far enough from the layer edge, the horizontal components of the ion velocity and the electric field are negligible. Therefore, the values of the surface tension, $s_{0,1}$, which determine the net force exerted upon the edge, are provided by the solution of the one-dimensional problem (1-3).

Before discussing the results of the numeric evaluation of the surface tension, it is convenient to introduce the normalized quantities. In the following, the coordinate, z , is measured in the units of the electron Debye length, $\lambda_D = \sqrt{T_e/4\pi e^2 n_0}$, the pressure is normalized to $n_0 T_e$, and the surface tension is normalized to $\lambda_D n_0 T_e$. The dimensionless surface density of




FIG. 2: The pressure (a) and the electric field (b) profiles in the sheath-presheath area evaluated for $\mu = 0.5$, $\eta = 0.1$. (a1) corresponds to the longitudinal pressure (9), (a2) corresponds to the transverse pressure (10), the dashed curve (a3) corresponds to the transverse pressure in the absence of dust ($\eta = 0$), (b1,b2) correspond to the electric field for $\eta \neq 0$ and $\eta = 0$. The vertical dashed lines in both figures show the position of the sheath-presheath interface ($z_s = 10.31$). Discontinuities at the curves (a1, a2, b1) are at the equilibrium position of the dust layer ($z_l = 3.94$).

the dust layer is $\eta = q\sigma/(2en_0\lambda_D)$ and the normalized grain mass is $\mu = Mge\lambda_D/(qT_e)$. Under the typical experimental conditions, $\mu \leq 1$, while $\eta \leq 0.1$. All computations discussed below have been performed with the sufficiently small value of the ionization frequency, $\nu_{ion}/\omega_{pi} = 0.01$, where ω_{pi} is the ion plasma frequency.

Examples of the pressure and the electric field profiles are depicted in Fig. 2. The net simulation length for this particular run is $z_0 = 68.5$. In order to make the effect discernible in the figure, we use here a sufficiently large value of the normalized dust density, $\eta = 0.1$. The longitudinal pressure drop (the curve 1 in Fig. 2a) at the charged layer is compensated by its weight; outside of the layer, the longitudinal pressure is constant. Contrary, the transverse pressure (10) is strongly inhomogeneous inside the sheath (the curves 2 and 3 in Fig. 2a). Of importance is that the dust reduces the transverse pressure between the layer and the electrode. As it is readily seen from Fig. 2b, the effect is mostly due to the reduction




FIG. 3: Dependence of the normalized surface tension on the dust density, η . 1 — $\mu = 0.8$, 2 — $\mu = 0.5$, 3 — $\mu = 0.1$, 4 — $\mu = 0.02$.




FIG. 4: The derivative of the surface tension with respect to the dust density versus the grain mass.

of the electric field.

The vertical dashed lines in Fig. 2 show the position of the sheath-presheath interface, z_s , for $\eta \neq 0$, which is conditionally defined as the point where the ion flow velocity is equal to the ion sound velocity. The dependence of the coordinate z_s on the dust parameters is rather weak. For this particular example, the dust layer levitates in the supersonic ion flow, $z_l < z_s$; with reducing grain weight, μ , it moves up to the subsonic area, $z_l > z_s$.

Fig. 3 shows the dependence of the surface tension on the dust layer density, η , for various values of the grain mass, μ . The curves 1-3 in Fig. 3 correspond to the dust layer levitating in the supersonic flow, $z_l < z_s$, the curve 4 corresponds to lighter dust grains situated in the presheath area, $z_l > z_s$. The depicted curves show that if the dust layer is located in the sheath, the surface tension increases with η , while with the dust layer shifted to the presheath, the surface tension may become a decreasing function of η . This is also shown in Fig. 4, where the dependence of the derivative $s'(\eta)|_{\eta=0}$ on the grain mass, μ , is plotted.

As we have already mentioned, the dependence of the surface tension on the dust density

determines the force acting upon the edge of the semi-infinite layer. If $s(\eta) > s(0)$ then the force $f_x = s(0) - s(\eta)$ is negative, as shown in Fig. 1, that is, the surface tension at the plasma-wall interface tends to expel the dust. Otherwise, the dust is drawn in the sheath. As it is seen from Figs. 3,4, in dependence on the grain mass and the dust density both alternatives are possible.

Here, we treat dust layer as a rigid incompressible medium. However, the short-range interaction between the dust grains result in the surface pressure, $p_s(\sigma)$, which also depends on the density [12]. The surface tension (11) reduces the surface pressure, that is, the net equation of state of the dust layer is given by $p = p_s(\sigma) - s(\sigma)$. Since $p_s(\sigma)$ exponentially tends to zero with decreasing σ , the compressibility, $dp/d\sigma$, of a rarefied dust layer is negative if $s'(0) > 0$. Such a medium is unstable. The detailed investigation of this instability and arising structures will be discussed elsewhere.

To summarize, charged dust may be treated as a surface-active substance that is capable of altering the surface tension at the plasma-electrode interface. Unlike usual surfactants, the dust increases the surface tension in a wide range of parameters. Although we did not discuss here the emergence of the two-dimensional void [3], it should be pointed out that a similar effect is well-known in liquids. To observe a void, it is enough to drop some liquid soap at the surface of water with floating fine powder.

This study was supported in part by the Netherlands Organization for Scientific Research (grant no. NWO 047.008.013) and the Russian Foundation for Basic Research (project no. 02-02-16439).

-
- [1] V.N. Tsytovich, G.E. Morfill, and H. Thomas, Plasma Phys. Rep. **28**, 675 (2002).
 - [2] P.K. Shukla and A.A. Mamun, *Introduction to Dusty Plasma Physics* (IOP Publishing, Bristol and Philadelphia, 2002).
 - [3] R. P. Dahiya *et al*, Phys. Rev. Lett. **89**, 125001 (2002).
 - [4] D. Samsonov and J. Goree, Phys. Rev. E **59**, 1047 (1999).
 - [5] H. Rothermel *et al*, Phys. Rev. Lett. **89**, 175001 (2002).
 - [6] G. E. Morfill *et al*, Phys. Rev. Lett. **83**, 1598 (1999).
 - [7] B.M. Annaratone *et al.*, Phys. Rev. E, **66**, 056411 (2002).

- [8] J. Goree, G. E. Morfill, V. N. Tsytovich, and S. V. Vladimirov, Phys. Rev. E **59**, 7055 (1999).
- [9] G. Morfill and V.N. Tsytovich, Phys. Plasmas **9**, 4 (2002).
- [10] K-U. Riemann, J. Phys. D **24**, 493 (1991).
- [11] S.V. Vladimirov and N.F. Cramer, Phys. Rev. E **62**, 2754 (2000).
- [12] G.A. Hebner, M.E. Riley, and K.E. Greenberg, Phys. Rev. E **66**, 046407 (2002).